

# ABDOMINAL PAIN STANDING ORDER

Initiate Immediate Supportive Care:

- Oxygen to maintain O2 sat  $\geq$  94%
- Complete primary and secondary survey as indicated
- Vital signs including FSBG and temperature as indicated

I  
N  
C  
L

Use standing order on patients with complaint of abdominal pain

E  
X  
C  
L

- Pregnant - follow [OB/GYN SO](#)
- Patients meeting [Trauma Triage Decision Scheme](#)
- Patients meeting [Critical Pediatric Triage Protocol](#)

O  
R  
D  
E  
R  
S

## BLS Care:

### Normal Volume Status

- Initiate IV NS/LR @ TKO (if permitted)
- Transport in position of comfort with supportive measures as indicated

### Volume Depleted

- Follow Normal Volume Status orders
- Bolus 20 ml/kg, reassess hemodynamic and pulmonary status at 500 ml intervals

## ALS Care:

- Follow BLS orders
- If angina equivalent present, consider cardiac monitor and 12 lead ECG
- Complaints of nausea and/or vomiting follow [Nausea/Vomiting/Diarrhea SO](#)
- Consider [Pain Management SO](#)

## Reference Note

### Clinical Signs of Hypoperfusion/Hypovolemia/ Hypotension

- Altered mental status
- Rapid pulse
- Hypoxia
- Cool Extremities
- Hypotension
- Prolonged cap refill
- Cyanosis
- Mottling

### Normal Vital Signs

Age	Heart Rate	Systolic BP
< 2 years of age	90-150	70 + (age in yrs x2)
2-3 years of age	80-140	70 + (age in yrs x2)
4-5 years of age	70-120	70 + (age in yrs x2)
6-9 years of age	60-115	70 + (age in yrs x2)
Over 10 years of age	60-110	$\geq$ 90